

100 學年度下學期數學系博士班資格考試  
(實變分析)

本試題卷共 2 頁，計 10 題計算證明題，每題 10 分，合計 100 分。

1. Prove the *Carathéodory theorem*: A set  $E$  is measurable if and only if for every set  $A$ ,

$$|A|_e = |A \cap E|_e + |A \setminus E|_e.$$

(Note:  $|A|_e$  denotes the outer measure of  $A$ .)

2. Prove that the set of points at which a sequence of measurable real-valued functions converges (to a finite limit) is measurable.
3. Let  $f$  be a function which is upper semi-continuous and finite on a compact set  $E$ . Show that if  $f$  is bounded above on  $E$ . Show also that  $f$  assumes its maximum on  $E$ , that is, that there exists  $x_0 \in E$  such that  $f(x_0) \geq f(x)$  for all  $x \in E$ .
4. Let  $f \in L(0, 1)$ . Show that  $x^k f(x) \in L(0, 1)$  for  $k = 1, 2, \dots$ , and  $\int_0^1 x^k f(x) dx \rightarrow 0$  as  $k \rightarrow \infty$ .
5. Let  $E$  be a measurable subset of  $\mathbb{R}^2$  such that for almost every  $x \in \mathbb{R}^1$ ,  $\{y \mid (x, y) \in E\}$  has  $\mathbb{R}^1$ -measure zero. Show that  $E$  has measure zero, and the for almost every  $y \in \mathbb{R}^1$ ,  $\{x \mid (x, y) \in E\}$  has measure zero.
6. (a) Write out the definition of the essential supremum  $\|f\|_\infty$  of a real-valued measurable function  $f$  on a measurable set  $E$ .
- (b) Let  $f$  be a real-valued measurable function on  $[0, 1]$ . Prove that  $\lim_{p \rightarrow \infty} \|f\|_p = \|f\|_\infty$ .
7. Let  $E$  be a measurable set in  $\mathbb{R}^n$ , and  $0 < p < q \leq \infty$ .
- (a) Prove that  $L^p(E) \cap L^\infty(E) \subset L^q(E)$ .
- (b) Prove that if  $|E| < \infty$ , then  $L^q(E) \subset L^p(E)$ .
8. Let  $f, g \in L^2(\mathbb{R}^n)$ . Prove that  $f + g \in L^2(\mathbb{R}^n)$  and  $\|f + g\|_2 \leq \|f\|_2 + \|g\|_2$ .

(背面尚有試題)

9. Let  $\{\varphi_k\}$  be an orthonormal system in  $L^2[0, 1]$ , and  $\{c_k\}$  be the Fourier series of a function  $f \in L^2[0, 1]$  with respect to the system  $\{\varphi_k\}$ .

(a) Prove that the Bessel's inequality  $\left(\sum_{k=1}^{\infty} |c_k|^2\right)^{1/2} \leq \|f\|_2$  holds.

(b) Find a necessary and sufficient condition so that the Parseval's identity  $\left(\sum_{k=1}^{\infty} |c_k|^2\right)^{1/2} = \|f\|_2$  holds, and prove your answer.

10. Let  $C[0, 1]$  denote the set of all real-valued continuous functions on  $[0, 1]$ , and the linear operator  $T : C[0, 1] \rightarrow \mathbb{R}$  be defined by  $T(f) = f(1)$  for all  $f \in C[0, 1]$ .

(a) Prove that  $T$  is a continuous linear functional on  $C[0, 1]$ .

(b) Prove that there exists an extension  $T^* : L^\infty[0, 1] \rightarrow \mathbb{R}^n$  of  $T$  such that  $T^*$  is a continuous linear functional on  $L^\infty[0, 1]$ , but there is no  $g \in L^1[0, 1]$  satisfying

$$T^*(f) = \int_{[0,1]} (f \times g) dx \quad \text{for all } f \in C[0, 1].$$

(試題結束)

101 學年度上學期數學系博士班資格考試  
(實變分析)

本試題卷共 2 頁，計 10 題計算證明題，每題 10 分，合計 100 分。

1. Let  $E$  be a measurable subset of  $\mathbb{R}$ , with  $|E| > 0$ . Prove that there exists a positive real number  $\varepsilon$  such that  $(-\varepsilon, \varepsilon) \subset E - E$ , where

$$E - E = \{x - y \mid x, y \in E\}.$$

2. Prove or disprove:

- (a) Any function  $f : [a, b] \rightarrow \mathbb{R}$  of bounded variation is measurable.  
(b) Any upper semicontinuous function  $f : [a, b] \rightarrow \mathbb{R}$  is measurable.

3. Let  $E$  be a measurable set in  $\mathbb{R}^n$  of finite measure. Prove that  $f : E \rightarrow \mathbb{R}$  is measurable if and only if for any  $\varepsilon > 0$ , there exists a closed subset  $F$  of  $E$  such that  $|E \setminus F| < \varepsilon$ , and  $f$  is continuous on  $F$ .

4. (a) State without proof the Egorov's theorem.

- (b) Let  $\langle f_k \rangle$  be a sequence of measurable functions on a measurable set  $E$  with  $|E| < \infty$ . If  $f_k$  converges to  $f$  a.e. in  $E$ , and  $\sup_k |f_k - f| \in L(E)$ , prove that  $\lim_{k \rightarrow \infty} \int_E f_k = \int_E f$ .

5. Let  $f : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$  satisfy for each  $x \in [0, 1]$ ,  $f(x, y)$  is a Lebesgue integrable function of  $y$ , and  $\frac{\partial f(x, y)}{\partial x}$  is a bounded function of  $(x, y)$ . Prove that  $\frac{\partial f(x, y)}{\partial x}$  is a measurable function of  $y$  for each  $x \in [0, 1]$ , and

$$\frac{d}{dx} \int_{[0,1]} f(x, y) dy = \int_{[0,1]} \frac{\partial f(x, y)}{\partial x} dy.$$

6. (a) State the definition for a finite function  $f$  on a finite interval  $[a, b]$  to be *absolutely continuous*.

- (b) Show that the function  $f(x) = x^\alpha$  is absolutely continuous on every bounded subinterval of  $[0, \infty)$  whenever  $\alpha > 0$ .

7. Let  $a_1, a_2, \dots, a_N$  be non-negative real numbers,  $p_1, p_2, \dots, p_N$  be positive real numbers with  $\sum_{j=1}^N (1/p_j) = 1$ . Show that

$$\prod_{j=1}^N a_j \leq \sum_{j=1}^N \frac{a_j}{p_j}.$$

(背面尚有試題)

8. Let  $\ell^\infty$  denote the normed linear space of all bounded real sequences. Is  $\ell^\infty$  separable? Justify your answer.
9. Suppose that  $f_k, f \in L^2$ , and that  $\int f_k g \rightarrow \int f g$  for all  $g \in L^2$ . If  $\|f_k\|_2 \rightarrow \|f\|_2$ , show that  $f_k \rightarrow f$  in  $L^2$  norm.
10. Let  $\Sigma$  be a  $\sigma$ -algebra on a set  $\mathcal{S}$ ,  $\{E_k\}$  be any sequence of sets in  $\Sigma$ , and  $\phi$  be a non-negative additive set function on  $\Sigma$ . Prove that

$$\phi\left(\liminf_{k \rightarrow \infty} E_k\right) \leq \liminf_{k \rightarrow \infty} \phi(E_k).$$

(試題結束)

103 學年度數學系博士班資格考試  
(實變分析)

※ 本試題卷共 8 題證明題

1. (a) Prove that every Borel measurable subset in  $\mathbb{R}^n$  is Lebesgue measurable.  
(b) Prove that there is a Lebesgue measurable subset in  $\mathbb{R}^n$  is not Borel measurable.  
(10%)
  
2. Prove or disprove (Please explain your answer):
  - (a) If  $f : [a, b] \rightarrow \mathbb{R}$  is a function of bounded variation, then  $f$  is Lebesgue measurable.
  - (b) If  $E$  is a Lebesgue measurable subset of  $\mathbb{R}$ , with  $|E| > 0$ , then there exist  $x, y \in E$  with  $x \neq y$  such that  $x - y$  is a rational number.
  - (c) If for each rational number  $a$ , the set  $\{x \in \mathbb{R}^n \mid f(x) > a\}$  is Lebesgue measurable, then  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is Lebesgue measurable.
  - (d) There exists a Riemann integrable function  $f : [0, 1] \rightarrow [0, 1]$  such that  $f$  is continuous at each rational point and discontinuous at each irrational point of  $[0, 1]$ .
  - (e) If  $f$  is Lebesgue integrable over  $E$ , then  $f$  is finite a.e. in  $E$ .  
(30%)
  
3. Prove that if  $f : [a, b] \rightarrow \mathbb{R}$  is a function of bounded variation, then  $f$  can be written as  $f = g + h$ , where  $g$  is absolutely continuous and  $h$  is singular, which are unique up to additive constants.  
(10%)
  
4. Prove that if  $f \in L^p(E)$  and  $f \geq 0$ , then  $\int_E f^p = p \int_0^\infty \alpha^{p-1} \omega(\alpha) d\alpha$ , where  $\omega$  is the distribution function of  $f$ , defined by  $\omega(\alpha) = |\{x \in E \mid f(x) > \alpha\}|$ .  
(10%)
  
5. Prove that if  $f \in L^p(\mathbb{R})$ , where  $1 \leq p < \infty$ , then for every  $\varepsilon > 0$  there is a continuous function  $g$  with compact support such that  $\|f - g\|_p < \varepsilon$ .  
(10%)
  
6. Prove that if  $f \in L(\mathbb{R}^n)$ , then the definite integral  $F(E) = \int_E f(x) dx$  is absolutely continuous with respect to Lebesgue measure.  
(10%)

7. For  $f, g \in L(\mathbb{R}^n)$ , we define the convolution of  $f$  and  $g$  by

$$(f * g)(x) = \int_{\mathbb{R}^n} f(x-y)g(y) dy \text{ for } x \in \mathbb{R}^n .$$

Prove that  $f * g \in L(\mathbb{R}^n)$ , and  $\|f * g\|_1 \leq \|f\|_1 \cdot \|g\|_1$ . (10%)

8. Let  $\{\varphi_k\}$  be an orthonormal system in  $L^2[0, 1]$ , and  $\{c_k\}$  be a sequence in  $\ell^2(\mathbb{R})$ . Prove that

there exists  $f \in L^2[0, 1]$  such that  $\sum_{k=1}^{\infty} c_k \varphi_k(x)$  is the Fourier series of  $f$  with respect to the orthonormal system  $\{\varphi_k\}$ . (10%)

103 學年度數學系博士班資格考試

(實變分析)

2015. 4. 30

※ 本試題卷共 8 題計算證明題

1. (a) Prove that if every measurable set  $E$  in  $\mathbb{R}^n$  can be expressed as  $E = F \cup Z$ , where  $F$  is a closed set and  $|Z| = 0$ .

(b) Let  $E_1$  and  $E_2$  be measurable subsets of  $\mathbb{R}^n$ . Prove that the product set  $E_1 \times E_2$  is a measurable subset of  $\mathbb{R}^n \times \mathbb{R}^n$ , and  $|E_1 \times E_2| = |E_1| \cdot |E_2|$ .

(15%)

2. Let  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  be measurable. Prove that the function  $g: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  defined by  $g(x, y) = f(x - y)$  is also measurable on  $\mathbb{R}^n \times \mathbb{R}^n$ .

(10%)

Hint : Show that there exists an invertible  $(2 \times 2)$  matrix  $A$  such that

$$\{ (x, y) \mid g(x, y) > a \} = A \left( \mathbb{R}^n \times \{ z \mid f(z) > a \} \right) \text{ for all } a \in \mathbb{R}.$$

3. Prove or disprove (Please explain your answer):

(a) There exists a Riemann integrable function  $f: [0, 1] \rightarrow [0, 1]$  such that  $f$  is continuous at each rational point and discontinuous at each irrational point of  $[0, 1]$ .

(b) There exists an increasing continuous function  $f$  whose derivative  $f'$  is Lebesgue integrable on  $[0, 1]$  such that  $\int_{[0, 1]} f' \neq f(1) - f(0)$ .

(10%)

4. (a) Prove carefully that for  $0 < a < b < \infty$ ,  $\int_{[0, \infty)} \int_{[a, b]} e^{-xy} \sin x \, dx \, dy = \int_{[a, b]} \frac{\sin x}{x} \, dx$ .

(b) Evaluate the Lebesgue integral  $\int_{(0, \infty)} \frac{\sin x}{x} \, dx$ .

(15%)

5. Let  $f: [0, 1] \rightarrow \mathbb{R}$  be measurable. Prove that if  $g(x, y) = f(x) - f(y)$  is Lebesgue integrable over  $[0, 1] \times [0, 1]$ , then  $f$  is Lebesgue integrable on  $[0, 1]$ .

(10%)

6. Let  $f_k : E \rightarrow \mathbb{R}$  be a sequence of measurable functions on  $E$ , where  $E$  is a measurable subset of  $\mathbb{R}^n$ , and  $1 \leq p < \infty$ .

(a) State the definition that  $\langle f_k \rangle$  converges to  $f$  in measure.

(b) State the definition that  $\langle f_k \rangle$  converges to  $f$  in  $L^p$ .

(c) Prove that if  $\langle f_k \rangle$  converges to  $f$  in  $L^p$ , then it converges to  $f$  in measure.

(15%)

7. (a) State without proof Holder inequality.

(b) Let  $E$  be a measurable subset of  $\mathbb{R}^n$ , with  $|E| \leq 1$ , and  $1 \leq p < q < \infty$ . Prove that for any measurable function  $f : E \rightarrow \mathbb{R}$ ,  $\|f\|_p \leq \|f\|_q$ .

(10%)

8. (a) Let  $f \in L^2(0, 1)$ . Prove that  $\lim_{k \rightarrow \infty} \int_0^{2\pi} f(x) \cos kx \, dx = \lim_{k \rightarrow \infty} \int_0^{2\pi} f(x) \sin kx \, dx = 0$ .

(b) Is (a) still true if  $f \in L^1(0, 1)$ ? Why?

(15%)



# 104 學年度數學系博士班資格考試

(實變分析)

2015. 10. 30

※ 本試題卷共六大題 (第一大題 50 分, 其餘各題每題 10 分)

1. Prove or disprove : (Please explain your answer)

(1) There is a Lebesgue measurable subset in  $\mathbb{R}^n$ , which is not Borel measurable.

(2) Any function  $f$  of bounded variation on  $[a, b]$  is Riemann integrable .

(3) There is a subset  $E$  of  $\mathbb{R}$ , with  $|E|_e > 0$ , satisfying for any  $x, y \in E$  with  $x \neq y$ ,  $x - y$  is not a rational number.

(4) There is a sequence  $\{E_k\}$  of disjoint sets such that  $\left| \bigcup_{k=1}^{\infty} E_k \right|_e < \sum_{k=1}^{\infty} |E_k|_e$ .

(5) If  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is Lebesgue measurable, then the function  $g : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  defined by  $g(x, y) = f(x - y)$  is also Lebesgue measurable on  $\mathbb{R}^n \times \mathbb{R}^n$ .

(6) Every Riemann integrable function  $f : [0, 1] \rightarrow \mathbb{R}$  is Lebesgue integrable.

(7) If  $f$  is Lebesgue integrable over  $E$ , then  $f$  is finite a.e. in  $E$ .

(8) If  $1 \leq p < q < \infty$ , then  $L^q[0, 1] \subset L^p[0, 1]$ .

(9) There exists an increasing continuous function  $f$  whose derivative  $f'$  is Lebesgue integrable on  $[0, 1]$  such that  $\int_{[0, 1]} f' \neq f(1) - f(0)$ .

(10) Any function  $f$  of bounded variation on  $[a, b]$  can be written as  $f = g + h$ , where  $g$  is absolutely continuous and  $h$  is singular.

(50%)

2. Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be an affine function defined by  $T(x) = Ax + u$ , where  $A$  is an  $n \times n$  matrix, and  $u$  is a fixed vector in  $\mathbb{R}^n$ . Prove that for any Lebesgue measurable set  $E$  of  $\mathbb{R}^n$ ,  $|T(E)| = |\det A| |E|$ .

(10%)

3. Let  $f : E \rightarrow \mathbb{R}$  be a Lebesgue measurable function, where  $E$  is a Lebesgue measurable

subset of  $\mathbb{R}^n$  with  $|E| < \infty$ . Prove that there exists a sequence  $\langle f_k \rangle$  of simple measurable functions on  $E$  such that  $\langle f_k \rangle$  converges almost uniformly to  $f$  in the following sense: for all  $\varepsilon > 0$ , there exists a closed subset  $F$  of  $E$  with  $|E \setminus F| < \varepsilon$ , such that  $\langle f_k \rangle$  converges uniformly to  $f$  on  $F$ . (Hint: You can apply Egorov Theorem) (10%)

4. Let  $f : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$  satisfy for each  $x \in [0, 1]$ ,  $f(x, y)$  is a Lebesgue integrable function of  $y$ , and  $\frac{\partial f(x, y)}{\partial x}$  is a bounded function of  $(x, y)$ . Prove that  $\frac{\partial f(x, y)}{\partial x}$  is a Lebesgue measurable function of  $y$  for each  $x \in [0, 1]$ , and

$$\frac{d}{dx} \int_{[0,1]} f(x, y) dy = \int_{[0,1]} \frac{\partial f(x, y)}{\partial x} dy. \quad (10\%)$$

5. Let  $f$  be nonnegative and Lebesgue measurable on a Lebesgue measurable subset  $E$  of  $\mathbb{R}^n$ . Prove that

$$\int_E f = \sup \sum_j [\inf_{x \in E_j} f(x)] |E_j|,$$

where the supremum is taken over all decompositions  $E = \cup_j E_j$  of  $E$  into the union of a finite number of disjoint Lebesgue measurable sets  $E_j$ . (10%)

6. Let  $\{\varphi_k\}$  be an orthonormal system in  $L^2[0, 1]$ , and  $\{c_k\}$  be a sequence in  $\ell^2(\mathbb{R})$ . Prove that there exists  $f \in L^2[0, 1]$  such that  $\sum_{k=1}^{\infty} c_k \varphi_k(x)$  is the Fourier series of  $f$  with respect to the orthonormal system  $\{\varphi_k\}$ . (10%)

# 105 學年度數學系博士班資格考試

(Real Analysis Qualifying Exam)

2016.10.31

1. Let  $E, F$  be measurable sets in  $\mathbb{R}^n$ ,  $B$  be a Borel set in  $[0, \infty)$ , and  $f : E \rightarrow [0, \infty)$  be a measurable function. Prove that the following 4 sets are measurable:

$$E \cup F, E \times F, f^{-1}\{B\}, \text{ and } R(f, E) = \{(x, y) \mid x \in E, 0 \leq y \leq f(x)\}. \quad (20\%)$$

2. (a) Use Caratheodory theorem to show that if  $E$  is a subset of  $\mathbb{R}^n$  satisfying the condition  $|G| = |G \cap E|_e + |G \cap E^C|_e$  for all open sets  $G$  in  $\mathbb{R}^n$ , then  $E$  is measurable.

- (b) If the condition in (a) is changed to  $|F| = |F \cap E|_e + |F \cap E^C|_e$  for all closed sets  $F$  in  $\mathbb{R}^n$ , is  $E$  measurable? Why? (10%)

3. Prove that if  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is a measurable function, then the function  $g : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ , defined by  $g(x, y) = f(2x - 3y)$ , is also measurable on  $\mathbb{R}^n \times \mathbb{R}^n$ . (10%)

(Hint: Find an invertible  $(2 \times 2)$  matrix  $A$  such that

$$\{(x, y) \mid g(x, y) > a\} = A\left(\mathbb{R}^n \times \{z \mid f(z) > a\}\right) \text{ for every } a \in \mathbb{R}.)$$

4. Let  $\langle f_k \rangle$  be a sequence of measurable functions on a measurable set  $E$  of  $\mathbb{R}^n$ .

(a) Use monotone convergence theorem to show that  $\int_E \sum_{k=1}^{\infty} |f_k| = \sum_{k=1}^{\infty} \int_E |f_k|$ .

- (b) Prove that if the series  $\sum_{k=1}^{\infty} \int_E |f_k|$  converges, then  $\sum_{k=1}^{\infty} f_k$  converges absolutely a.e. in

$$E, \text{ and } \sum_{k=1}^{\infty} \int_E f_k = \int_E \sum_{k=1}^{\infty} f_k. \quad (16\%)$$

5. (a) Prove that if  $f \in L(E)$ , then for all  $\varepsilon > 0$ , there is  $\delta > 0$  such that  $\int_A |f| < \varepsilon$  for all measurable subsets  $A$  of  $E$  with  $|A| < \delta$ .

- (b) Use Egoroff theorem to show that if  $\langle f_k \rangle$  is a sequence of measurable functions that converges to  $f$  a.e. in  $E$ , with  $|E| < \infty$ , and  $\sup_k |f_k - f| \in L(E)$ , then  $\lim_{k \rightarrow \infty} \int_E f_k = \int_E f$ .

- (c) Use Tonelli theorem to show that if  $f, g \in L(\mathbb{R}^n)$ , then  $\int_{\mathbb{R}^n} |f(x-y) \times g(y)| dy < \infty$  for a.e.  $x \in \mathbb{R}^n$ . (24%)

6. Let  $\{\varphi_k\}$  be an orthonormal system in  $L^2[0, 1]$ . Prove that  $\{\varphi_k\}$  is complete if, and only if,

Parseval's formula  $\|f\| = \left( \sum_{k=1}^{\infty} |c_k|^2 \right)^{1/2}$  holds for every  $f \in L^2[0, 1]$ , where the numbers  $c_k$  are the Fourier coefficients of  $f$  with respect to the system  $\{\varphi_k\}$ . (10%)

7. Use Radon-Nikodym theorem to show that for any continuous linear functional  $T$  on

$L^2[0, 1]$ , there exists a unique function  $g \in L^2[0, 1]$  such that  $T(f) = \int_{[0,1]} f \times g$  for every  $f \in L^2[0, 1]$ . (10%)

# 106 學年度數學系博士班資格考試

(Real Analysis Qualifying Exam)

2017.10.31

\*\*\*Each problem is worth 10 points.\*\*\*

1. Determine which function is Riemann (improper) integrable on  $E$ ? Lebesgue integrable on  $E$ ? Explain your answer.

$$f(x) = \begin{cases} 1, & \text{if } x \in [0,1] \cap \mathbb{Q} \\ x, & \text{if } x \in [0,1] \cap \mathbb{Q}^c \end{cases} \text{ on } E = [0,1] \text{ and } g(x) = \frac{\sin x}{x} \text{ on } E = [1, \infty).$$

2. Prove that (Caratheodory Theorem) a subset  $E$  in  $\mathbb{R}^n$  is measurable if and only if for every set  $A$  in  $\mathbb{R}^n$ ,  $|A|_e = |A \cap E|_e + |A \setminus E|_e$ .

3. Construct a sequence of disjoint sets  $E_1, E_2, E_3, \dots$  in  $\mathbb{R}$  such that  $\left| \bigcup_{k=1}^{\infty} E_k \right|_e \neq \sum_{k=1}^{\infty} |E_k|_e$ .

4. Prove that there exists a Lebesgue measurable set in  $\mathbb{R}$ , which is not a Borel set.

5. Prove that if  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  is measurable, then the function  $g: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ , defined by  $g(x, y) = f(x + 2y)$ , is also measurable on  $\mathbb{R}^n \times \mathbb{R}^n$ .

6. Let  $\langle f_k \rangle$  be a sequence of measurable functions on a measurable set  $E$  of  $\mathbb{R}^n$ . Prove that if the series  $\sum_{k=1}^{\infty} \int_E |f_k|$  converges, then  $\sum_{k=1}^{\infty} f_k$  converges absolutely *a.e.* in  $E$ , and

$$\sum_{k=1}^{\infty} \int_E f_k = \int_E \sum_{k=1}^{\infty} f_k.$$

7. Suppose that  $f \in L(\mathbb{R})$  and  $\iint_{\mathbb{R}^2} f(3x)f(x+2y) dx dy = 1$ , calculate  $\int_{\mathbb{R}} f(x) dx$ .

8. (a) Prove that if  $f: [a, b] \rightarrow \mathbb{R}$  is bounded, Lebesgue integrable, and  $F(x) = \int_{[a,x]} f$ ,

then  $F$  is absolutely continuous, and  $F' = f$  *a.e.* in  $[a, b]$ .

(b) Is (a) still true, if  $f$  is unbounded? Why?

9. Let  $f \in L^p(\mathbb{R}^n)$ ,  $1 < p, q < \infty$ , and  $\frac{1}{p} + \frac{1}{q} = 1$ . Prove that  $\|f\|_p = \sup_{\|g\|_q \leq 1} \left| \int_{\mathbb{R}^n} f(x) \times g(x) dx \right|$ .

10. (a) Let  $f \in L^2(0, 2\pi)$ . Prove that  $\lim_{k \rightarrow \infty} \int_0^{2\pi} f(x) \cos kx dx = \lim_{k \rightarrow \infty} \int_0^{2\pi} f(x) \sin kx dx = 0$ .

(b) Is (a) still true, if  $f \in L^1(0, 2\pi)$ ? Why?



# 108 學年度數學系博士班資格考試(實變分析)

## Real Analysis Qualifying Exam

2019.10.31

1. It is known from Caratheodory theorem that a subset  $E$  of  $\mathbb{R}^n$  is measurable if and only if  $|A| = |A \cap E|_e + |A \setminus E|_e$  for all sets  $A$  in  $\mathbb{R}^n$ . Prove or disprove :
- (a) If  $|G| = |G \cap E|_e + |G \setminus E|_e$  for all open sets  $G$  in  $\mathbb{R}^n$ , then  $E$  is measurable.
- (b) If  $|F| = |F \cap E|_e + |F \setminus E|_e$  for all closed sets  $F$  in  $\mathbb{R}^n$ , then  $E$  is measurable. (12%)
2. (a) Let  $f : [0, 1] \rightarrow \mathbb{R}$  be a continuous function and  $B$  denote the Borel  $\sigma$ -algebra in  $\mathbb{R}$ . Prove that the family  $\Gamma = \{E \subset \mathbb{R} \mid f^{-1}(E) \text{ is measurable}\}$  is a  $\sigma$ -algebra containing  $B$ .
- (b) Prove that there exists a measurable subset of  $[0, 1]$ , but not a Borel set. (12%)
3. (a) Prove that every linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  maps measurable subsets of  $\mathbb{R}^n$  into measurable sets.
- (b) Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a measurable function, and  $a, b \in \mathbb{R}$ . Prove that the function  $g : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ , defined by  $g(x, y) = f(ax + by)$ , is also measurable on  $\mathbb{R}^n \times \mathbb{R}^n$ . (12%)
4. Prove that if  $f : \mathbb{R} \rightarrow \mathbb{R}$  is a measurable function satisfying  $f(x + y) = f(x) + f(y)$  for all  $x, y \in \mathbb{R}$ , then  $f$  must be linear. (10%)
5. (a) Prove that if  $f \in L(E)$ , then  $f$  is finite *a.e.* in  $E$ .
- (b) Suppose that  $\langle f_k \rangle$  is a sequence of measurable functions on a measurable subset  $E$  of  $\mathbb{R}^n$ , and  $\sum_{k=1}^{\infty} \int_E |f_k|$  converges. Prove that  $\sum_{k=1}^{\infty} f_k$  converges absolutely *a.e.* in  $E$ , and  $\sum_{k=1}^{\infty} \int_E f_k = \int_E \sum_{k=1}^{\infty} f_k$ . (12%)
6. Let  $\langle f_k \rangle$  be a sequence of measurable functions on a measurable subset  $E$  of  $\mathbb{R}^n$ , with  $|E| < \infty$ , and  $|f_k(x)| \leq M_x < \infty$  for all  $k$  and for each  $x \in E$ . Prove that for all  $\varepsilon > 0$ , there is a closed subset  $F$  of  $E$  and a positive number  $M$  such that  $|E \setminus F| < \varepsilon$  and  $|f_k(x)| \leq M$  for all  $k$  and for all  $x \in F$ . **(Hint : You can apply Lusin theorem)** (10%)

7. Use Tonelli theorem to show that if  $f : E \rightarrow [0, \infty)$  is a measurable function on a measurable subset  $E$  of  $\mathbb{R}^n$ , and  $\omega(\alpha) = |\{x \in E \mid f(x) > \alpha\}|$ , then  $\int_E f = \int_0^\infty \omega(\alpha) d\alpha$ .

(Hint :  $\int_E f = \iint_{R(f,E)} 1 dx dy$ , where  $R(f,E) = \{(x,y) \mid x \in E, 0 \leq f(x) \leq y\}$ .) (10%)

8. Let  $f : [0,1] \times [0,1] \rightarrow \mathbb{R}$  be a measurable function. Prove that if the iterated integral

$\int_{[0,1]} \int_{[0,1]} |f(x,y)| dx dy$  exists and is finite, then  $f \in L([0,1] \times [0,1])$ , and

$$\iint_{[0,1] \times [0,1]} f = \int_{[0,1]} \int_{[0,1]} f(x,y) dx dy = \int_{[0,1]} \int_{[0,1]} f(x,y) dy dx. \quad (10\%)$$

9. Let  $\{\varphi_k\}$  be any orthonormal basis for  $L^2(E)$  over  $\mathbb{R}$ .

(a) Prove that  $\{\varphi_k\}$  must be countable and complete.

(b) Prove that any function  $f \in L^2(E)$  satisfies Parseval formula with respect to  $\{\varphi_k\}$ ;

that is,  $\|f\|_2 = \left( \sum_{k=1}^{\infty} |c_k|^2 \right)^{\frac{1}{2}}$ , where  $\{c_k\}$  is the sequence of Fourier coefficients of  $f$ .

(12%)

109 學年度數學系博士班資格考試(實變分析)

Real Analysis Qualifying Exam

2021.4.28

1. Let  $f(x) = \begin{cases} 0, & \text{if } x \in [0,1] \\ 1, & \text{if } x \in (1,2] \end{cases}$ ,  $\alpha(x) = \begin{cases} 0, & \text{if } x \in [0,1] \\ 1, & \text{if } x \in [1,2] \end{cases}$ , and  $\beta(x) = \begin{cases} x, & \text{if } x \in [0,1] \\ x^2, & \text{if } x \in [1,2] \end{cases}$ .
- (a) Is  $f$  Riemann-Stieltjes integrable to  $\alpha$  on  $[0,2]$ ? Why?
- (b) Is  $f$  Riemann-Stieltjes integrable to  $\beta$  on  $[0,2]$ ? Why? (12%)
2. (a) Let  $f : [0,1] \times [0,1] \rightarrow \mathbb{R}$  be a measurable function and  $B$  be a Borel set in  $\mathbb{R}$ . Prove that  $f^{-1}(B)$  is measurable in  $[0,1] \times [0,1]$ .
- (b) Let  $f$  and  $g$  be measurable on  $[0,1]$ . Prove that the function  $F : [0,1] \times [0,1] \rightarrow \mathbb{R}$ , defined by  $F(x, y) = f(x) \times g(y)$ , is measurable on  $[0,1] \times [0,1]$ . (12%)
3. Let  $f : E \rightarrow \mathbb{R}$  be a measurable function on a measurable subset  $E$  of  $\mathbb{R}^n$ . Prove that for all  $\varepsilon > 0$ , there is a Borel set  $B$  in  $E$ , with  $|E \setminus B| < \varepsilon$ , and a sequence  $\langle f_k \rangle$  of Borel measurable functions such that  $\langle f_k(x) \rangle$  converges increasingly to  $|f(x)|$  for all  $x \in B$ . (10%)
4. Let  $\langle f_k \rangle$  be a sequence of measurable functions on a measurable subset  $E$  of  $\mathbb{R}^n$ , and  $\sum_{k=1}^{\infty} \int_E |f_k|$  converges. Prove that  $\sum_{k=1}^{\infty} |f_k|$  converges *a.e.* in  $E$ , and  $\sum_{k=1}^{\infty} \int_E f_k = \int_E \sum_{k=1}^{\infty} f_k$ . (10%)
5. Let  $\langle f_k \rangle$  be a sequence of increasing functions on  $[a, b]$ , and  $\sum_{k=1}^{\infty} f_k(x)$  converge to  $f(x)$  for each  $x \in [a, b]$ . Prove that  $\sum_{k=1}^{\infty} f'_k(x)$  converges to  $f'(x)$  for *a.e.*  $x$  in  $E$ . (10%)



6. Let  $f : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$  satisfy that for each  $x \in [0, 1]$ ,  $f(x, y)$  is a Lebesgue integrable function of  $y$ , and  $\frac{\partial f(x, y)}{\partial x}$  is a bounded function of  $(x, y)$ . Prove that  $\frac{\partial f(x, y)}{\partial x}$  is a measurable function of  $y$  for each  $x \in [0, 1]$ , and

$$\frac{d}{dx} \int_{[0,1]} f(x, y) dy = \int_{[0,1]} \frac{\partial f(x, y)}{\partial x} dy. \quad (10\%)$$

7. Let  $E$  be a measurable subset of  $\mathbb{R}^n$ . Prove that  $f : E \rightarrow \mathbb{R}$  is measurable if and only if the region  $R(f, E)$  is measurable, where  $R(f, E) = \{(x, y) \mid x \in E, 0 \leq f(x) \leq y\}$ .
- (12%)

8. (a) Let  $f$  be measurable on  $E$ , and  $1 < p < q < \infty$ , with  $\frac{1}{p} + \frac{1}{q} = 1$ . Prove that

$$\int_E |fg| \leq \left( \int_E |f|^p \right)^{\frac{1}{p}} \left( \int_E |f|^q \right)^{\frac{1}{q}}$$

- (b) Let  $f$  be measurable on  $E$  with  $0 < |E| < \infty$ , and  $1 \leq p < q < \infty$ . Prove that

$$\left( \frac{1}{|E|} \int_E |f|^p \right)^{\frac{1}{p}} \leq \left( \frac{1}{|E|} \int_E |f|^q \right)^{\frac{1}{q}}. \quad (12\%)$$

9. Define the operator  $T : C[0,1] \rightarrow \mathbb{R}$  by  $T(f) = f(1)$  for all  $f \in C[0,1]$ , where  $C[0,1]$  denotes the Banach space of all real-valued continuous functions on  $[0, 1]$ .

- (a) Prove that  $T$  is a continuous linear functional on  $C[0,1]$ .

- (b) Prove that there exists a continuous linear functional  $T^* : L^\infty[0,1] \rightarrow \mathbb{R}$  such that

$$T^*(f) = T(f) \text{ for all } f \in C[0,1], \text{ but there exists no function } g \in L^1[0,1] \text{ satisfying } T^*(f) = \int_{[0,1]} (f \times g) dx \text{ for all } f \in C[0,1]. \quad (12\%)$$